861-01-1991

PORC

Brewer's Blackbird - Jan 12, 1991

This single bird woo seen about 2:30 on 1/12/410m a fresh manure strip near McAlevey'st Ft, Huntingdom 6. I observed it from my of at a distance of 30-40 yards. There was bright sunshine at a 9.0 angle (on my right) from my line of vision to the bird.

The bird had a very light white a yellow eyes Phus
It had to be grackle, rusty blackbird, on Brewar's.

I ruled oct common grackle due to size. The bird had a short tail (in companion to grackles). I woo able to compare its length directly with starlings and horned lanks in the field in my binocular at the same time as I observed this bird. It's length was very similar to starling and slightly longer than horned lanks.

I ruled out rusty blackbird based on plumage. This bird was black, not "rusty". On the lugat scalight, it's head had a distinct purple sheen and its body had a green-blue sheen.

The bird kept mostly to itself although starlings, horned lanks, mourning doves rock dove and I water pipet were also on the manure strip

I observed the bird for 25 minutes through 8.5 × 44 Swift Audoban binoculars

Y

Greg Grove BE1, Box 483 Petersbung 16669 861-01-1991

Brewer's Blackbird Huntingdon County Greg Grove

This single bird was seen about 2:30 on 12 January p.m. 1991 on a fresh manure strip near McAlevy's Fort, Huntingdon County. I observed it from my car at distance of 30-40 There vards. bright sunshine at a angle (on my right) from my line of vision to the bird.

The bird had very light white or yellow eyes. Thus it had to be a Common Grackle (Quiscalus quiscula), Rusty Blackbird (Euphagus carolinus), or Brewer's Blackbird (Euphagus).

ruled Ι grackle due to size. The bird had a short tail (in comparison to grackles). I was able to compare its length directly with European Starlings (Sturnus vulgaris) and Horned Larks (Eremophila alpestris) in field in my binoculars at the same time as I bird. observed this Its length was verv similar to starlings and slightly larger than the larks.

I ruled out Rusty Blackbird based on plumage. This bird was black, not "rusty." In the bright sunlight its head had a distinct purple sheen and its body had a green-blue sheen.

The bird kept mostly to itself,

although starlings, Horned Larks, Mourning Doves (Zenaida macroura), Rock Doves (Columba livia), and one American Pipit (Anthus rubescens) were also on the manure strip.

I observed the bird for 25 minutes through 8.5X44 Swift Audubon binoculars.

RR 1, Box 483 Petersburg, Pennsylvania 16669

Pennsylvania Ornithological Records Committee

Voting Tabulation - Round # of

Species:

Brewer's Blackbird

Date of Sighting:

January 12, 1991

Observer(s):

Greg Grove

Date of Submission:

1991 Huntinsday

Submitted by:

Greg Grove

Member	Class I	Class II	Class III	Class IV	Class V		
					A	В	С
E. Kwater			X				
B. Haas			\times				
F. Haas			X				
R. Leberman			X				
G. McWilliams			X				
S. Santner			X				
P. Schwalbe			X				
TOTALS			7				
DECISION			X				

Comments:

Signature (Secretary):

3m Haas

Date: //-6-9/